

Blow-up results for some second order hyperbolic inequalities with a nonlinear term with respect to the velocity

M. Jazar and R. Kiwan
Lebanese University and Université de Tours

February 7, 2008

Abstract

We give sufficient conditions on the initial data so that a semilinear wave inequality blows-up in finite time. Our method is based on the study of an associated second order differential inequality. The same method is applied to some semilinear systems of mixed type.

AMS Subject Classifications: Primary 35L70, 34B40, 34D05; secondary 34A40

Keywords: Wave equations and systems, Hyperbolic-elliptic systems, Hyperbolic-parabolic systems, Ordinary differential equations and inequalities, Finite time blow-up, Asymptotic behavior of solutions, Invariant regions

1 Introduction

Let Ω a bounded regular¹ domain of \mathbb{R}^N and let us denote respectively by $|\cdot|$ and (\cdot, \cdot) the natural norm and inner product of the space $H = L^2(\Omega)$. Let $V \hookrightarrow H$ be a real Hilbert space that is continuously and densely embedded into H . Denote by $\|\cdot\|$ and $\langle \cdot, \cdot \rangle_V$ respectively the norm and the inner product of V and by $L \in \mathcal{L}(V, V^*)$ the unique linear continuous operator from V to V^* , the dual of V , satisfying

$$\langle Lu, v \rangle_{V^*, V} = \langle u, v \rangle_V \quad (u, v) \in V \times V.$$

Moreover we assume that the operator L admits an eigenvalue λ (that we may assume to be positive without loss of generality) with a positive associated eigenfunction $\varphi \in V$.

¹The regularity needed here is that ensuring the existence of a first eigenvalue with positive associated eigenfunction for L .

Consider the semilinear evolution second order partial differential inequality

$$\begin{cases} u'' + Lu \geq g(u') & \text{in } \mathcal{D}^*((0, T) \times \Omega), \\ u(0, x) = u_0(x) & \text{on } \Omega \\ u_t(0, x) = u_1(x) & \text{on } \Omega, \end{cases} \quad (1)$$

on the phase space $(u_0, u_1) \in V \times H$.

The model problem is a semilinear wave equation with Dirichlet boundary conditions, that is $V = H_0^1(\Omega)$, $L = -\Delta$, see the book of S. Alinhac [1] for a short view on semilinear hyperbolic problems and the question of blow-up. The particular case of equation with $g(x) = -x|x|^r$ is a dissipative semilinear wave equation that has been extensively studied in the literature, and many results on the existence of global solutions as well as the decay estimates have been established. See for instance [5, 4, 10, 18].

The case of equation with $g(x) = x|x|^r$ is more complicated. For instance, a local existence-uniqueness of solutions is only guaranteed for small r and more regular initial data. See for instance [5, 7] for equations with nonlinearities on u and u_t . In [9], A. Haraux showed that this problem (with $g(x) = x|x|^x$) has no nontrivial global and bounded solutions [9, Proposition 1.2]. Moreover, he constructed examples of blowing-up solutions with arbitrarily small initial data [9, Proposition 1.4 and Corollary 1.5]. See also the work of H. Levin, S. R. Park and J. Serrin [13] and the references therein for a study of the long-time behavior of solutions for equations with $f(u) - g(u_t)$ on the right hand side of the equation on $\Omega = \mathbb{R}^N$.

In this paper we consider Problem (1) where g is a positive convex function $C^1(\mathbb{R}, \mathbb{R}^+)$ satisfying the growth condition

$$g(x) \geq C|x|^q \quad \text{for all } x \in \mathbb{R}, \quad (2)$$

where $C > 0$ is a constant and $q > 1$ if the dimension $N = 1$ or 2 or $1 < q \leq \frac{N+2}{N-2}$ if $N \geq 3$. We provide sufficient conditions on the initial data (u_0, u_1) so that the solution of Problem (1) blows-up in finite time.

The method used here is the so-called “eigenfunction method” which leads to the study of a second order ordinary differential inequality that we make using invariant regions. This is the purpose of the second section. In the third section we apply this to the semilinear wave equation (1). **S. Kaplan [11] was the first to use Fourier coefficient’s method in this setting. Later, R. Glassey [8] applied this method to nonlinear wave equations. In particular he treated an equation similar to (1) in the whole space \mathbb{R}^N . H. Levine [12, Section 5] considered equation similar to (1) on bounded domain with Dirichlet boundary conditions. Under some hypotheses on the nonlinearity g , the eigenvalue λ and the initial data, Levine gives a necessary condition for finite time blow-up. Our assumptions on the nonlinearity are different, in particular g does not need to be nondecreasing.**

Moreover, the method we use for the study of the ODI is different from that given in [12]. However, the model case, the case of equality with $g(s) = C|s|^q$, is covered by Levine's work (see [12, Section 5]), but our assumptions on the initial data are weaker. In the third Section (Remark 3) we give a comparison between Levine's result and ours.

In the fourth section we apply our result to semilinear wave systems of the form:

$$\begin{cases} u_{tt} - \Delta u & \geq |v_t|^p \\ v_{tt} - \Delta v & \geq |u_t|^q, \end{cases} \quad (3)$$

by means of an extension of results of Section 2 to an associated system of second order ordinary differential inequalities. In the last section we obtain similar results for systems of mixed type: hyperbolic-elliptic and hyperbolic-parabolic systems (see for instance the work of Pohozaev and Véron [14]).

Acknowledgment. The authors thank the anonymous referee of this paper for his report. He pointed out to us the papers of R. Glassey [8] and H. Levine [12].

2 Finite time blow-up for a second order ordinary differential inequality

Consider the ODI:

$$\begin{cases} v'' + av \geq bv'^q, & t \geq 0 \\ v(0) = v_0, & v'(0) = v_1, \end{cases} \quad (4)$$

where a, b are positive constants.

Notice that in [2, 17] one can find a complete study of the equation: $u'' + a|u|^{p-1}u = b|u'|^{q-1}u'$ where $a, b > 0$ and $p, q > 1$. In [15, 16], Ph. Souplet proved the existence of some global solutions to equation (4).

Concerning the ODI (4) we distinguish two cases: $q \leq 2$ and $q > 2$

Proposition 1 *Let $q \leq 2$, $\alpha = \frac{2a}{bq} \left(\frac{4a}{b^2q} \right)^{\frac{2-q}{2q-2}}$ and $v_0, v_1 \in \mathbb{R}$ satisfying*

$$\left| \begin{array}{l} 0 < v_1, \\ av_0 + \alpha < \frac{b}{2}v_1^q \quad \text{if } v_1 < \left[\frac{4a}{b^2q} \right]^{\frac{1}{2q-2}}, \\ av_0 < \frac{b}{2}v_1^q \quad \text{if } v_1 \geq \left[\frac{4a}{b^2q} \right]^{\frac{1}{2q-2}}. \end{array} \right. \quad (5)$$

Then all solutions of (4) blow-up in finite time. Moreover,

$$v'(t) \geq \left[v_1^{1-q} - \frac{q-1}{2}bt \right]^{\frac{1}{1-q}}. \quad (6)$$

The proof of the proposition is based on the idea that the inequality $v'' \geq bv'^q$ provide blowing up solution in finite time $T < \frac{v_1^{1-q}}{b(q-1)}$. We look then for invariant regions under the ODI (4) in order to obtain $av^p \leq b\varepsilon v'^q$.

Proof of Proposition 1. We show first that the region \mathcal{D} defined by (5) is invariant under (4). Denote by F the function defined over \mathbb{R} by

$$F(x) := \begin{cases} 0 & \text{for } x \leq x_1 := -\frac{\alpha}{a} \\ \left[\frac{2}{b}(ax + \alpha)\right]^{\frac{1}{q}} & \text{for } x_1 \leq x \leq 0 \\ \left[\frac{4a}{b^2q}\right]^{\frac{1}{2q-2}} & \text{for } 0 \leq x \leq x_2 := \frac{b}{2a} \left[\frac{4a}{b^2q}\right]^{\frac{q}{2q-2}} \\ \left[\frac{2a}{b}x\right]^{\frac{1}{q}} & \text{for } x \geq x_2. \end{cases}$$

The region \mathcal{D} is then defined by $[y > F(x)]$.

Let v be a solution of (4) satisfying (5). Setting $x = v$ and $y = v_t$, the ODI (4) can be transformed into the dynamical system:

$$\begin{cases} x' &= P(x, y) := y, \\ y' &= Q(x, y) \geq by^q - ax, \\ x(0) &= v_0, \\ y(0) &= v_1. \end{cases} \quad (7)$$

In order to prove that the region \mathcal{D} is invariant we will show that the vector field defining the dynamical system is “entering” along the curve $y = F(x)$. This is clear on the semi-axis $(y'O)$ and on the other segment. On the arc $x > x_2$,

$$\frac{y'}{x'} \geq \frac{by^q - ax}{y} \geq \frac{b}{2}y^{q-1} \geq \frac{2a}{bqy^{q-1}} = F'(x).$$

Now, on the arc $x_1 < x < 0$ we have

$$\frac{y'}{x'} \geq \frac{\alpha}{y} \geq F'(x).$$

Therefore for all $t > 0$ we have $\frac{b}{2}v_t^q \geq av$, and then $v'' > \frac{1}{2}bv'^q$. Integrating we get (6). \square

The following proposition gives a similar result for the case $q > 2$. we discuss later the difference between the corresponding invariant regions in these two cases.

Proposition 2 *Let $q > 2$ and $v_0, v_1 \in \mathbb{R}$ satisfying*

$$\begin{cases} 0 < v_1, \\ bqv_1^{q-2}(bv_1^q - av_0) - a > 0. \end{cases} \quad (8)$$

Then all solutions of (4) blow-up in finite time. Moreover,

$$v'(t) \geq \left[v_1^{1-q} - (q-1)(1-\varepsilon)bt \right]^{\frac{1}{1-q}} \quad (9)$$

where $\varepsilon \in (0, 1)$ is a constant satisfying:

$$\varepsilon bq v_1^{q-2}(\varepsilon b v_1^q - a v_0) - a \geq 0. \quad (10)$$

Proof. First note that, using (8) and taking

$$f(\varepsilon) = \varepsilon bq v_1^{q-2}(\varepsilon b v_1^q - a v_0) - a,$$

there exists at least an $\varepsilon \in (0, 1)$ satisfying $f(\varepsilon) > 0$.


Setting $A = \varepsilon b v_1^q - a v_0$, we have $v_1^{q-2} \geq \frac{a}{\varepsilon bq A}$. Denote by \mathcal{D}' the domain $[y > F_2(x)]$, where $F_2(x) := \left[\frac{1}{\varepsilon b(ax+A)} \right]^{\frac{1}{q}}$. Since y' remains positive, as long as (x, y) belongs to \mathcal{D}' , then for all $t > 0$, $(y(t))^{q-2} \geq \frac{a}{\varepsilon bq A}$. Following the same arguments as in the last proposition, we have, along the curve $y = F_2(x)$,

$$\frac{y'}{x'} \geq \frac{by^q - ax}{y} \geq \frac{\varepsilon by^q - ax}{y} = \frac{A}{y} \geq \frac{a}{\varepsilon bq y^{q-1}} = F_2'(x).$$


□

Remark 1 *(On the admissible initial data region)*

In the following figures we draw the curve limiting the region for two values of q . The admissible region in each case is colored in gray.



$a = 1, b = 2$ and $q = 1.5$.



$a = b = 1$ and $q = 2.5$.

Remark 2 *The same method works for the differential inequality*

$$v'' + av^p \geq bv'^q,$$

where $p \leq 1$.

3 Blow-up criteria for a semilinear wave inequality

In this section we assume that we have a Sobolev injection type theorem: $V \hookrightarrow L^{2^*}(\Omega)$, with $2^* = \frac{2N}{N-2}$. Consider the problem

$$\begin{cases} u \in C([0, T]; V) \cap C^1([0, T]; H) \\ u'' + Lu \geq g(u') & \text{in } \mathcal{D}^*((0, T) \times \Omega), \\ u(0, x) = u_0(x) & \text{on } \Omega \\ u_t(0, x) = u_1(x) & \text{on } \Omega, \end{cases} \quad (11)$$

where g is a positive convex function $C^1(\mathbb{R}, \mathbb{R}^+)$ satisfying the growth condition (2), $q > 1$ if the dimension $N = 1$ or 2 or $1 < q \leq \frac{N+2}{N-2}$ if $N \geq 3$ and $(u_0, u_1) \in V \times H$. Indeed, local existence is guaranteed under hypotheses of this type on both the power q and the smoothness of the initial data, see for instance [6, 9]. Under such hypotheses, local existence can be easily obtained by classical fixed point argument and abstract semi-group theory.

Let λ be the first positive eigenvalue of the operator L on H , and φ a nonnegative associated eigenfunction satisfying $\int_{\Omega} \varphi = 1$. Denote also by $v_i := \int_{\Omega} u_i \varphi$, $i = 0, 1$. As for the ODI (4), we distinguish 2 cases depending on the value of q .

Theorem 1 *Let $(u_0, u_1) \in V \times H$ such that v_0 and v_1 satisfy:*

$$\begin{aligned} \text{if } q \leq 2 \quad & \left| \begin{array}{l} 0 < v_1, \\ \lambda v_0 + \alpha < \frac{C}{2} v_1^q \quad \text{if } v_1 < \left(\frac{4\lambda}{C^2 q}\right)^{\frac{1}{2q-2}}, \\ \lambda v_0 \leq \frac{C}{2} v_1^q \quad \text{if } v_1 \geq \left(\frac{4\lambda}{C^2 q}\right)^{\frac{1}{2q-2}}. \end{array} \right. \\ \text{and if } q > 2 \quad & \left| \begin{array}{l} 0 < v_1, \\ Cq v_1^{q-2} (Cv_1^q - \lambda v_0) - \lambda > 0. \end{array} \right. \end{aligned}$$

where $\alpha := \frac{2\lambda}{Cq} \left(\frac{4\lambda}{C^2 q}\right)^{\frac{2-q}{2q-2}}$. Then every solution of (11) blows-up in finite time

$$T_{max} \leq T^* := \frac{v_1^{1-q}}{(1-\varepsilon)C(q-1)},$$

where $\varepsilon := \frac{1}{2}$ if $q \leq 2$, and if $q > 2$, ε satisfy $\varepsilon Cq v_1^{q-2} (Cv_1^q - \lambda v_0) - \lambda \geq 0$. Moreover,

$$\|u_t(t)\|_{L^1(\Omega)} \geq \|\varphi\|_{L^\infty(\Omega)}^{-1} \left[v_1^{1-q} - (q-1)(1-\varepsilon)Ct \right]^{\frac{1}{1-q}}.$$

Proof. Set $v(t) := \int_{\Omega} u(t)\varphi dx$. By the elliptic regularity theory $\varphi \in V$ and then v is twice differentiable with $v' = \int u_t \varphi$ and $v'' = \langle u_{tt}, \varphi \rangle_{V^*, V}$ that we write also as $\int_{\Omega} u_{tt} \varphi dx$.

Multiply the equation (11) by $\varphi \geq 0$ and then integrate over Ω , we get after integrating by parts

$$\begin{aligned} \int_{\Omega} u_{tt} \varphi dx - \int_{\Omega} Lu \varphi dx &\geq C \int_{\Omega} |u_t|^q \varphi dx \\ v''(t) + \lambda v(t) &\geq C |v'(t)|^q, \end{aligned}$$

where we used Jensen's lemma. Using Propositions 1 and 2 we deduce that v blows-up before the time T^* and

$$\begin{aligned} \|\varphi\|_{L^\infty(\Omega)} \|u_t(t)\|_{L^1(\Omega)} &\geq \left| \int_{\Omega} u_t(t) \varphi dx \right| = |v'(t)| \\ &\geq \left[v_1^{1-q} - C(q-1)(1-\epsilon)t \right]^{\frac{-1}{q-1}}. \end{aligned}$$


□

Remark 3 (Comparison with the results of Levine [12])

In [12, Theorem 5.1], Levine shows a blow-up result for a class of equations. Levine's result can be easily applied to equation (11) with $g(s) = C|s|^q$. However, the assumptions he needs on the initial data are not the same. Indeed, by [12, Hypothesis (5.5)], v_0 and v_1 satisfy

$$v_1 > v_0 > s_0$$

for some s_0 (s_0 should be in this case greater than $(\frac{\lambda+1}{C})^{\frac{1}{q-1}}$). This gives a region of the form



to be compared with the regions of Remark 1.

Remark 4 According to Remark 2, the same method could be generalized to inequalities of the form ($p \leq 1$)

$$u_{tt} - \Delta u^p \geq g(u_t).$$

4 Application to systems of semilinear wave inequalities

In this section we show how to extend the same method to the following system

$$\left\{ \begin{array}{l} u, v \in C([0, T]; V) \cap C^1([0, T]; H) \\ u'' + Lu \geq |v'|^p \\ v'' + Lv \geq |u'|^q \end{array} \right\} \quad \text{in } \mathcal{D}^*((0, T) \times \Omega), \quad (12)$$

$$\left\{ \begin{array}{ll} u(0, x) = u_0(x), & u_t(0, x) = u_1(x) \\ v(0, x) = v_0(x), & v_t(0, x) = v_1(x) \end{array} \right. \quad \text{on } \Omega,$$

where (u_0, u_1) and (v_0, v_1) are in $V \times H$ and the powers p and q are greater than 1 if the dimension $N = 1$ or 2 or in $\left(1, \frac{N+2}{N-2}\right]$ if $N \geq 3$. For simplicity we choose a simple form of the non linearity, although a general form is possible. For this, we introduce the following system of ODI:

$$\left\{ \begin{array}{ll} U'' + aU \geq V'^p & t \geq 0 \\ V'' + aV \geq U'^q & t \geq 0 \\ U(0) = U_0, & U'(0) = U_1 \\ V(0) = V_0, & V'(0) = V_1. \end{array} \right. \quad (13)$$

The method of Section 2 could be easily extended to the previous system as follows:

Proposition 3 *Let $1 < p \leq q$, U_0, V_0, U_1 and V_1 satisfying*

$$\left| \begin{array}{ll} U_1, V_1 > 1, & U_0 V_0 \geq U_1 V_1, \\ U_1^q & \geq [a + \frac{1}{p}] V_0, \\ V_1^p & \geq [a + \frac{1}{p}] U_0. \end{array} \right. \quad (14)$$

Then every solution (U, V) of (13) blows-up in finite time. Moreover we have

$$U'(t) + V'(t) \geq \left[(U_1 + V_1)^{1-p} - \frac{p-1}{1+ap} 2^{1-p} t \right]^{\frac{1}{1-p}}.$$

Proof. Setting $x = U$, $y = V'$, $z = V$ and $t = U'$, then the system (13) is transformed into the 4-d dynamical system

$$\left\{ \begin{array}{ll} x' = t, & x(0) = U_0, \\ t' \geq y^p - ax, & t(0) = U_1, \\ z' = y, & z(0) = V_0, \\ y' \geq t^q - az, & y(0) = V_1. \end{array} \right.$$

Set $\alpha := 1 + \frac{1}{ap}$ and $f_p(x) := H(x)(\alpha ax)^{\frac{1}{p}}$, where H is the Heaviside function. Denote by $\mathcal{D}_1 := [y \geq f_p(x)]$ and $\mathcal{D}_2 := [t \geq f_q(z)]$. The hypothesis (14) can

then be read as $(U_0, U_1) \in \mathcal{D}_1$ and $(V_0, V_1) \in \mathcal{D}_2$.

First notice that as long as (x, y, z, t) remains in $\mathcal{D}_1 \times \mathcal{D}_2$, U'' and V'' remain positive, so for all $t > 0$, $t^{q-1}y^{p-1} \geq U_1^{q-1}V_1p - 1 = \frac{U_1^q V_1^p}{U_1 V_1} \geq \alpha^2 a^2 \frac{U_0 V_0}{U_1 V_1} \geq \alpha^2 a^2$.

In order to show that (x, y, z, t) could not exit the domain $\mathcal{D}_1 \times \mathcal{D}_2$ let's examine the vector field along the boundary. On $[t = f_q(z)]$ we have

$$\begin{aligned} \frac{y'}{x'} &\geq \frac{t^q - az}{t} = \frac{(\alpha - 1)az}{t} = \frac{\alpha - 1}{\alpha} t^{q-1} \\ &\geq \frac{\alpha - 1}{\alpha} \frac{\alpha^2 a^2}{y^{p-1}} = ap(\alpha - 1)f'_p(x) = f'_p(x). \end{aligned}$$

A similar calculation holds for the other boundary.

Finally, setting $\beta := \frac{\alpha-1}{\alpha}$, we have:

$$\begin{cases} U'' &\geq \beta V'^p, \\ V'' &\geq \beta U'^q. \end{cases}$$

Adding these two equations, using the fact that $U' \geq 1$ and $V' \geq 1$ for all $t > 0$, we get:

$$\frac{1}{2\beta} W'' \geq \frac{1}{2} (V'^p + U'^q) \geq \frac{1}{2} (V'^p + U'^p) \geq 2^{-p} W'^p, \quad (15)$$

where $W := U + V$. Integrating we obtain:

$$W'(t) \geq \left(W_1^{1-p} - (p-1)\beta 2^{1-p} t \right)^{\frac{1}{1-p}}.$$

□

In order to apply the last result to the system (12), denote by $U_i := \int_{\Omega} u_i \varphi$ and $g_i := \int_{\Omega} v_i \varphi$, $i = 0, 1$. Using the same method as in the proof of Theorem 1 and applying the Proposition 3 we get directly:

Theorem 2 *Let $1 < p \leq q$, (u_0, u_1) and (v_0, v_1) in $V \times H$ such that U_0, V_0, U_1 and V_1 satisfy the hypothesis*

$$\left| \begin{array}{ll} U_1, V_1 > 1 & U_0 V_0 \geq U_1 V_1, \\ U_1^q & \geq [\lambda + \frac{1}{p}] V_0, \\ V_1^p & \geq [\lambda + \frac{1}{p}] U_0. \end{array} \right. \quad (16)$$

Then every solution of (12) blows-up in finite time. Moreover, we have

$$\|u_t + v_t\|_{L^1(\Omega)} \geq \|\varphi\|_{L^\infty(\Omega)}^{-1} \left[(U_1 + V_1)^{1-p} - \frac{p-1}{1+\lambda p} 2^{1-p} t \right]^{\frac{1}{1-p}}.$$

5 Application to systems of mixed types

In this section we show how to apply the result of section 3 to some hyperbolic-elliptic and hyperbolic-parabolic systems.

5.1 Hyperbolic-elliptic system

Consider the following system

$$\left\{ \begin{array}{l} u, v \in C([0, T]; H_0^1(\Omega)) \cap C^1([0, T]; L^2(\Omega)) \\ u_{tt} - \Delta u \geq |v_t|^q \\ -\Delta v = u \end{array} \right\} \quad \text{in } \mathcal{D}^*((0, T) \times \Omega), \quad (17)$$

$$\left\{ \begin{array}{ll} u(0, x) = u_0(x), & u_t(0, x) = u_1(x) \\ v(0, x) = v_0(x) \end{array} \right. \quad \begin{array}{l} \text{on } \Omega \\ \text{on } \Omega, \end{array}$$

where $q > 1$.

Theorem 3 *Let $u_0, v_0 \in H_0^1(\Omega)$ and $u_1 \in L^2(\Omega)$ such that $U_0 := \int_{\Omega} u_0 \varphi dx$ and $U_1 := \int_{\Omega} u_1 \varphi dx$ satisfy:*

$$\text{if } q < 2 \quad \left| \begin{array}{l} 0 < U_1, \\ \lambda U_0 + \alpha < \frac{\lambda^{-q}}{2} U_1^q \quad \text{if } U_1 < \left(\frac{4\lambda^{2q+1}}{q} \right)^{\frac{1}{2q-2}}, \\ \lambda U_0 \leq \frac{\lambda^{-q}}{2} U_1^q \quad \text{if } U_1 \geq \left(\frac{4\lambda^{2q+1}}{q} \right)^{\frac{1}{2q-2}}. \end{array} \right.$$

$$\text{and if } q > 2 \quad \left| \begin{array}{l} 0 < U_1, \\ q\lambda^{-q} U_1^{q-2} (\lambda^{-q} U_1^q - \lambda U_0) - \lambda > 0, \end{array} \right.$$

where $\alpha := \frac{2\lambda^{1+q}}{q} \left(\frac{4\lambda^{2q+1}}{q} \right)^{\frac{2-q}{2q-2}}$. Then every solution of (17) blows-up in finite time

$$T_{max} \leq T^* := \frac{U_1^{1-q} \lambda^q}{(1-\varepsilon)(q-1)},$$

where $\varepsilon := \frac{1}{2}$ if $q \leq 2$, and if $q > 2$, ε satisfy $\varepsilon C q v_1^{q-2} (\varepsilon C v_1^q - \lambda v_0) - \lambda \geq 0$. Moreover,

$$\|u_t(t)\|_{L^1(\Omega)} \geq \|\varphi\|_{L^\infty(\Omega)}^{-1} \left[U_1^{1-q} - (q-1)(1-\varepsilon)\lambda^{-q}t \right]^{\frac{1}{1-q}}.$$

Proof. Denote by $U := \int_{\Omega} u \varphi dx$ and $V := \int_{\Omega} v \varphi dx$. Differentiate the second equation of (17), multiply by φ , then integrate over Ω we get

$$\lambda V'(t) = U'(t). \quad (18)$$

Multiplying the first equation of (17) by φ and then integrating over Ω we get, after using (18),

$$U''(t) + \lambda U(t) \geq \lambda^{-q} |U'(t)|^q.$$

We conclude applying the propositions 1 and 2 with $a = \lambda$ and $b = \lambda^{-q}$. \square

5.2 Hyperbolic-parabolic system

Consider the following system

$$\left\{ \begin{array}{l} u, v \in C([0, T]; H_0^1(\Omega)) \cap C^1([0, T]; L^2(\Omega)) \\ u_{tt} - \Delta u \geq |v_t|^q \\ (u - v)_t - \Delta(u - v)^m \leq \beta(u - v)^p \\ u - v \geq 0 \\ u(0, x) = u_0(x), \quad u_t(0, x) = u_1(x) \\ v(0, x) = v_0(x) \end{array} \right\} \quad \begin{array}{l} \text{in } \mathcal{D}^*((0, T) \times \Omega), \\ \text{on } [0, T] \times \Omega, \\ \text{on } \Omega \\ \text{on } \Omega, \end{array} \quad (19)$$

where $q > 1$, $m \geq 1 \geq p$ and $\beta \in \mathbb{R}$.

Theorem 4 *Let $u_0, v_0 \in H_0^1(\Omega)$ and $u_1 \in L^2(\Omega)$. Set $U_0 := \int_{\Omega} u_0 \varphi dx$, $V_0 := \int_{\Omega} v_0 \varphi dx$ and $U_1 := \int_{\Omega} u_1 \varphi dx$. Assume that $\beta \leq 0$ or $[\beta \leq \lambda \text{ and } U_0 - V_0 \geq 1]$, $u_0 - v_0 \geq 0$ and*

$$\text{if } q < 2 \quad \left| \begin{array}{l} 0 < U_1, \\ \lambda U_0 + \alpha < \frac{1}{2} U_1^q \quad \text{if } U_1 < \left(\frac{4\lambda}{q}\right)^{\frac{1}{2q-2}}, \\ \lambda U_0 \leq \frac{1}{2} U_1^q \quad \text{if } U_1 \geq \left(\frac{4\lambda}{q}\right)^{\frac{1}{2q-2}}. \end{array} \right. \quad (20)$$

$$\text{and if } q > 2 \quad \left| \begin{array}{l} 0 < U_1, \\ q U_1^{q-2} (U_1^q - \lambda v_0) - \lambda > 0, \end{array} \right.$$

where $\alpha = \frac{2\lambda}{q} \left(\frac{4a}{q}\right)^{\frac{2-q}{2q-2}}$.

Then every solution of (19) blows-up in finite time

$$T_{max} \leq T^* := \frac{U_1^{1-q}}{(1-\varepsilon)(q-1)},$$

where $\varepsilon := \frac{1}{2}$ if $q \leq 2$, and if $q > 2$, ε satisfy $\varepsilon q v_1^{q-2} (\varepsilon v_1^q - \lambda v_0) - \lambda \geq 0$. Moreover,

$$\begin{aligned} \|v_t(t)\|_{L^1(\Omega)} &\geq \|u_t(t)\|_{L^1(\Omega)} \\ &\geq \|\varphi\|_{L^\infty(\Omega)}^{-1} \left[U_1^{1-q} - (q-1)(1-\varepsilon)t \right]^{\frac{1}{1-q}}. \end{aligned}$$

Proof. Denote by $U := \int_{\Omega} u \varphi dx$ and $V := \int_{\Omega} v \varphi dx$. Multiplying the second equation of (19) by φ and then integrating over Ω we get after using Jensen's lemma

$$(U - V)' \leq \int_{\Omega} [\beta(u - v)^p - \lambda(u - v)^m] \varphi \quad (21)$$

$$\leq \beta(U - V)^p - \lambda(U - V)^m. \quad (22)$$

By the hypothesis on β and using the diffusion property, $U' \leq V'$. Set

$$T := \max \{0 \leq t < T_{max} \text{ s.t. } U'(s) \geq 0 \quad \forall s \in [0, t]\}.$$

Since $U_1 > 0$, one has $T > 0$. For all $t \in [0, T)$ we have $0 \leq U' \leq V'$ and, hence, $0 \leq U'^q \leq V'^q$. Multiplying the first equation of (19) by φ then integrating over Ω we get

$$U'' + \lambda U \geq U'^q \quad (23)$$

for all $t \in [0, T)$. In order to apply Propositions 1 and 2 we need to prove that T is large enough. Indeed, by the hypothesis (20), (U, U') remains in the invariant region defined by (20) (see the proof of Propositions 1 and 2). Thus $U'(t) > 0$ for all t and we obtain the result by applying Propositions 1 and 2 since (23) holds for all t . \square

References

- [1] S. Alinhac. Blowup for nonlinear hyperbolic equations. Progress in Nonlinear Differential Equations and their Applications, 17. Birkhäuser Boston, Inc., Boston, MA, 1995.
- [2] M. Balabane; M. Jazar and Ph. Souplet. Oscillatory blow-up in nonlinear second order ODE's: the critical case. Discrete Contin. Dyn. Syst. **9** (2003), no. 3, 577–584.
- [3] M. F. Bidaut-Véron and M. Bouhar. On characterization of solutions of some nonlinear differential equations and applications. SIAM J. Math. Anal. **25** (1994), no. 3, 859–875.
- [4] A. Carpio. Existence of global solutions to some nonlinear dissipative wave equations. J. Math. Pures Appl. (9) **73** (1994), no. 5, 471–488.
- [5] Th. Cazenave. Uniform Estimates for Solutions of Nonlinear Klein-Gordon Equations. J. Funct. Analysis **60** (1985), 36–55.
- [6] Th. Cazenave and A. Haraux. An introduction to semilinear evolution equations. Oxford Lecture Ser. Math. Appl. 13, Oxford 1998.
- [7] V. Georgiev and G. Todorova. Existence of a solution of the wave equation with nonlinear damping and source terms, J. Differential Equations **109** (1994), 295–308.
- [8] R. T. Glassey. Blow-up theorems for nonlinear wave equations. Math. Z. **132** (1973), 183–203.
- [9] A. Haraux. Remarks on the wave equation with a nonlinear term with respect to the velocity. Portugal. Math. **49** (1992), no. 4, 447–454.
- [10] A. Haraux and E. Zuazua. Decay estimates for some semilinear damped hyperbolic problems. Arch. Rational Mech. Anal. **100** (1988), no. 2, 191–206.
- [11] S. Kaplan. On the growth of solutions of quasilinear parabolic equations. Comm. Pur. Appl. Math. **16** (1957), 305–330.
- [12] H. A. Levine. Nonexistence of global weak solutions to some properly and improperly posed problems of mathematical physics: the method of unbounded Fourier coefficients. Math. Ann. **214** (1975), 205–220.
- [13] H. A. Levine and S. R. Park and J. Serrin. Global Existence and Global Nonexistence of Solutions of the Cauchy Problem for a Nonlinearly Damped Wave Equation. Journ. Math. Anal. Appl., **228** (1998), pp. 181–205.

- [14] S. Pohozaev and L. Véron. Blow-up results for nonlinear hyperbolic inequalities. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, vol. XXIX (2000), 393–420.
- [15] Ph. Souplet. Etude des solutions globales de certaines équations différentielles ordinaires du second ordre non linéaires, *C. R. Acad. Sci. Paris*, t. **313**, Série 1 (1991), 365–370.
- [16] Ph. Souplet. Existence of exceptional growing-up solutions for a class of nonlinear second order ordinary differential equations, *Asymptotic Analysis*, **11** (1995), 185–207.
- [17] Ph. Souplet. Critical exponents, special large-time behavior and oscillatory blow-up in nonlinear ODE's. *Differential Integral Equations* **11** (1998), no. 1, 147–167.
- [18] Ph. Souplet. Sharp stability estimates for quasi-autonomous evolution equations of hyperbolic type. *Quart. Appl. Math.* **57** (1999), no. 1, 55–85.

M. Jazar
 Lebanese University
 Mathematics Department
 P.O.Box 155-012
 Beirut Lebanon
 mjazar@ul.edu.lb

R. Kiwan
 Université de Tours
 Laboratoire de Mathématiques
 et Physique Théorique
 UMR 6083 du CNRS
 Parc de Grandmont
 37200 Tours France
 kiwan@lmpt.univ-tours.fr